



Ref : CERMA2010/E1

Analgesic and anti-inflammatory effects of *Cassia siamea* Lam. stem bark extracts.

G.F. Nsonde Ntandou^{a,b,c}, J.T. Banzouzi^{b,d}, B. Mbatchi^{a,b}, R.D.G. Elion-Itou^a, A.W. Etou-Ossibi^a, S. Ramos^d, F. Benoit-Vical^{e,f,g}, A.A. Abena^a and J.M. Ouamba^h

a Laboratoire de Biochimie et Pharmacologie, Faculté des Sciences de la Santé, Université Marien NGOUABI, Brazzaville, B.P. 69, Congo

b Centre d'Etude et de Recherche Médecins d'Afrique (CERMA), B.P. 45, Brazzaville, Congo

c Laboratoire de Pharmacologie, Centre d'Etudes sur les Ressources Végétales (CERVE), B.P. 1249, Brazzaville, Congo

d Institut de Chimie des Substances Naturelles (ICSN-CNRS), 1 Avenue de la Terrasse-Bat 27, 91198 Gif-sur-Yvette Cedex, France

e Service de Parasitologie-Mycologie, Centre Hospitalier Universitaire de Rangueil, Université de Toulouse et Faculté de Médecine de Rangueil, Université de Toulouse III, UPS, TSA 50032, 31059 Toulouse Cedex 9, France

f CNRS, LCC (Laboratoire de Chimie de Coordination) UPR8241, 205, route de Narbonne, F-31077 Toulouse, France

g Université de Toulouse III, UPS, LCC, 118, route de Narbonne, F-31077 Toulouse, France

h Unité de Chimie du Végétal et de la Vie, Faculté des Sciences, Université Marien Nguabi, Brazzaville, B.P. 69, Congo

Journal of Ethnopharmacology Volume 127, vol 1, 108-111 (2010)

Abstract

AIM OF THE STUDY: The present study was carried out to investigate analgesic and anti-inflammatory activities of *Cassia siamea* Lam stem bark extracts. We have also determined the cytotoxicity of each extract.

MATERIALS AND METHODS: *C. siamea*, a widespread medicinal plant traditionally used in sub-Saharan Africa, was collected in Congo Brazzaville. Stem bark was extracted with petroleum ether (CSE1), chloroform (CSE2), ethanol (CSE3) and water (CSE4). Analgesic, anti-inflammatory and antipyretic activities of these extracts were assessed in rats with hot plate test, paw pressure and carrageenan induced paw oedema. Cytotoxicity was assessed against KB and Vero cells.

RESULTS: At the doses used (100, 200, and 400mg/kg) ethanol and water extracts showed significant and dose-dependent analgesic and anti-inflammatory effects. None of the extracts had cytotoxic activity on KB and Vero cell lines and the most active extracts (CSE3 and CSE4) had no acute toxicity.

CONCLUSIONS: The study highlighted the analgesic and anti-inflammatory of *C. siamea* stem bark. Four major families of compounds present in the plant may explain these activities: triterpenes (lupeol, oleanolic acid, ursolic acid, friedelin, betulin), flavonoids (apigenin, kaempferol, luteolin), anthraquinones (emodin), phytosterols (stigmasterol, beta-sitosterol).

Keywords:

Cassia siamea; *Fabaceae*; Stem bark extracts; Analgesic; Anti-inflammatory; Antipyretic